




Diabetes in Abbeville County, 2012



Produced by the South Carolina Department of Health and Environmental Control

Impact of Diabetes in the United States


- Diabetes affects 25.8 million Americans and contributes to approximately 231,000 deaths a year.
- Diabetes is becoming more common in the United States. From 1980 through 2011, the number of Americans with diagnosed diabetes has more than tripled (from 5.6 million to 20.9 million).
- The direct and indirect costs of diabetes in the U.S. were nearly \$174 billion in 2008.
- 69,701 people died of diabetes in the U.S. in 2010.

Diabetes in South Carolina

- Approximately 421,549 adults in South Carolina are diagnosed with diabetes. African Americans have a 42% higher prevalence than Whites.
- The total direct costs of hospitalizations and emergency room visits were over \$4.7 billion in 2012.
- Diabetes is the seventh leading cause of death in South Carolina, claiming 1,186 lives in 2012


Behavioral Risk Factors

- Major behavioral risk factors, such as obesity and physical inactivity are partially responsible for development of type 2 diabetes (Fig 1).



Diabetes Prevalence


- An estimated 11.2% of adults in Region 1 suffer annually from diabetes (Fig 2).



*County-level BRFSS not available for 2012.

Hospital and Emergency Room Discharges

- In 2012, there were 48 hospitalizations for diabetes in Abbeville County for all ages.
- There were an additional 501 hospital discharges with diabetes-related conditions in Abbeville County.
- In 2012, there were 92 emergency room (ER) visits for diabetes as the primary diagnosis, among which 57 (62%) were by African-American patients.
- There were 1348 ER visits for diabetes as a related condition.




- Medicare paid for 67% of the ER and Hospitalization charges in Abbeville County (Fig 3).

Complications

- 43% of patients hospitalized in SC for diabetes had complications in 2012; 38% of hospitalizations in Abbeville County (Table 1).
- In 2011, more than 74% of people in SC with diabetes had hypertension, compared with only 30% of people without diabetes.


diabetes Complication	Hospital Admissions	Percent of diabetes Hospital Admissions	Total charges
Renal Failure	161	30%	\$5,800,200
Lower Extremity Amputation	18	3%	\$876,900
Ketoacidosis	15	3%	\$781,300
Retinopathy	10	2%	\$503,900

- In 2011, more than 67% of people with diabetes had high cholesterol.
- 2012, the rates for lower extremity amputations among blacks in SC were 3times the rates among whites.




Mortality

- diabetes is the sixth leading cause of death in Abbeville County.
- A total of 10 people in Abbeville County died from diabetes in 2012.



- The Abbeville County mortality rate was 29.9/100,000 in 2012, below the state average.
- Abbeville County ranks #23 in diabetes mortality in South Carolina in 2010-2012.

Fig. 6 Diabetes Mortality 2010-2012



For More Information about diabetes contact:

Division of Diabetes, Heart Disease, Obesity and School Health
 DHEC Bureau of Community Health and Chronic Disease Prevention
<http://www.dhec.sc.gov/health/chcdp/diabetes/index.htm>
 (803) 898-1635

Diabetes Initiative of South Carolina
<http://www.musc.edu/diabetes/>



Generated by: Division of Chronic Disease Epidemiology, Bureau of Community Health and

http://www.scdhec.gov/hs/epidata/state_reports.htm

Data Sources:

Behavioral Risk Factors and Diabetes Prevalence:
 SC BRFSS
<http://www.scdhec.gov/administration/phsis/biostatistics/brfss/>

Mortality Data:
 Vital Statistics, DHEC Division of Biostatistics 803-898-3627



Hospital Discharges and ER Visits:
 Office of Research and Statistics SC Budget and Control Board (803) 898-9940



www.orss.state.sc.us
 For more details, please view the Technical Document below
www.scdhec.gov/health/epidata/docs/EpiTechNotes.pdf