Flu Watch

South Carolina Department of Health and Environmental Control Division of Acute Disease Epidemiology

Week Ending August 20, 2016 (MMWR Week 33) All data are provisional and may change as more reports are received.

In this issue:	
Summary	2
I. Confirmatory testing	3
II. Positive rapid tests	6
III. ILINet	8
IV. Hospitalizations and deaths	10
V. National influenza surveillance	12
VI. SC influenza surveillance components	13
VII. Definitions for influenza surveillance	14

MMWR Week 33 at a Glance:

Influenza Activity Synopsis:

During MMWR week 33 influenza activity in South Carolina increased; however, remained below historic baseline levels. South Carolina reported **No Activity.**

Laboratory surveillance:

- 14 laboratory-confirmed cases of influenza was reported from 6 counties.
- Of the positive specimens reported this season, 22,282 (66.1%) are influenza A, 10,792 (32.0%) are influenza B, 504 (1.5%) are influenza A/B, and 145 (0.4%) are influenza unknown subtype.

ILI Activity (South Carolina baseline is 2.05%):

 The percentage of visits to sentinel providers for influenzalike illness (0.17%) was below South Carolina's baseline.
ILI percentages represent ILI activity reported by less than half of enrolled sentinel providers. Therefore, ILI percentages may not be representative of actual flu activity.

Hospitalizations:

 No laboratory confirmed influenza-associated hospitalizations were reported. Since October 4, 2015, 1,856 laboratory confirmed influenza associated hospitalizations have been reported.

Deaths:

• No laboratory confirmed influenza-associated deaths were reported. Since October 4, 2015 forty-four (44) laboratory confirmed influenza associated deaths have been reported.


Summary of Laboratory Confirmed Tests, ILI Activity, Influenza Associated Hospitalizations and Deaths Compared to Previous Week and Previous Season


			-			
	Current week	Previous week	Change from previous week	<i>Cumulative</i> (2015-16)	<i>Cumulative</i> (2014-15)	Cumulative change 2015-16 compared to 2014-15
Number of positive confirmatory tests (culture, RT-PCR, DFA, IFA)	0	0		1,099	1,191	▼ 8%
Positive rapid antigen tests	14	1		32,624	53,990	V 40%
Percent of ILI visits reported by ILINet providers	0.17%	0.00%	A 0.17%			
Number of lab confirmed flu hospitalizations	0	0		1,856	3,358	V 45%
Number of lab confirmed flu deaths	0	0		44	155	72%

Positive confirmatory influenza test results* Current MMWR Week (8/14/16 – 8/20/16)					
	BOL and reference labs				
Number of positive confirmatory tests	0				
Influenza A unsubtyped					
Influenza A H1N1					
Influenza A H3N2					
Influenza B					
Other					
Includes culture, RT-PCR	, DFA, and IFA				

For the current MMWR reporting week, 0 positive confirmatory tests was reported.

<i>Positive confirmatory influenza test results*</i> <i>Cumulative (10/4/15 – 8/20/16)</i>				
BOL and reference labs				
1099				
167				
590				
79				
262				
1				







Map of all Laboratory Confirmed Cases (n) and Population Case Raters/100,000 by County October 4, 2015 - August 20, 2016


II. Positive Rapid Antigen Tests

During the most recent MMWR week, 14 positive rapid antigen tests were reported. Of these, 9 were influenza A, 4 were influenza B, and 1 was other/unknown. This compares to 6 during this same week last year.






During the most recent MMWR week, 0.17%* of patient visits to SC ILINet providers were due to ILI. This is below the state baseline (2.05%). This ILI percentage compares to 0.56% this time last year. Reports were received from providers in 4 counties, representing 2 of the 4 regions. The statewide percentage of ER visits with fever-flu syndrome was 3.92%.



ILI percentage is dependent upon the number of reporting providers and can be greatly influenced by a single provider with high numbers of ILI.



The SC fever flu percentage only includes data from hospitals emergency departments and urgent care centers participating in SC syndromic surveillance.

Influenza-Like Illness Reported by Sentinel Providers August 14, 2016 – August 20, 2016

County	ILI %	ILI % County	
Abbeville		Greenwood	NR
Aiken	NR	Hampton	
Allendale		Horry	
Anderson	NR	Jasper	
Bamberg		Kershaw	
Barnwell		Lancaster	
Beaufort	NR	Laurens	NR
Berkeley	NR	Lee	
Calhoun		Lexington	NR
Charleston	NR	Marion	
Cherokee		Marlboro	
Chester		McCormick	
Chesterfield		Newberry	
Clarendon		Oconee	
Colleton		Orangeburg	
Darlington		Pickens	
Dillon		Richland	
Dorchester	NR	Saluda	
Edgefield		Spartanburg	
Fairfield		Sumter	
Florence	NR	Union	
Georgetown	NR	Williamsburg -	
Greenville	NR	York 0.00	

NR: No reports received ---: No enrolled providers


For the current MMWR reporting week, no laboratory confirmed influenza-associated hospitalizations were reported by 3 hospitals. No laboratory confirmed influenza-associated deaths were reported. Since October 4, 2015, 1,856 laboratory confirmed influenza-associated hospitalizations and 44 laboratory confirmed influenza-associated deaths have been reported. Laboratory confirmation for hospitalizations and deaths includes culture, PCR, DFA, IFA, and rapid antigen detection test.


	_	Current M	IMWR Wee	ek (8/14/16	- 8/20/1	6)	
	0-4	5-17	18-49	50-64	65+	Unknown	Total
Hospitalizations							0
Deaths							0

Cumulative (10/4/15 – 8/20/16)							
	0-4	5-17	18-49	50-64	65+	Unknown	Total
Hospitalizations	188	128	447	476	614	3	1856
Deaths	0	1	8	17	18		44

* Lab confirmation for hospitalizations and deaths includes culture, PCR, DFA, IFA, and rapid tests.

Influenza associated deaths by Region				
Region	Total			
Lowcountry	12			
Midlands	7			
Pee Dee	11			
Upstate	14			





V. National surveillance (8/7/16 - 8/13/16)

CDC has begun summer publication of a much-abbreviated Flu View report. To view national data, go to: http://www.cdc.gov/flu/weekly/

VI. South Carolina Influenza Surveillance Components

South Carolina influenza surveillance consists of mandatory and voluntary reporting systems for year-round influenza surveillance. These networks provide information on influenza virus strain and subtype and influenza disease burden.

Mandatory Reporting

Positive confirmatory test reporting

Positive influenza culture, PCR, DFA, and IFA results from commercial laboratories must be reported to DHEC within 3 days electronically via SCION or using a DHEC 1129 card.

Positive rapid antigen test reporting

Summary numbers of positive rapid influenza tests and influenza type identified must be sent to the regional health department by fax or email before noon on Monday for the preceding week.

Influenza deaths

All laboratory confirmed influenza deaths (adult and pediatric) must be reported to DHEC within 24 hours. These include results from viral culture, PCR, rapid flu tests, DFA, IFA or autopsy results consistent with influenza.

Influenza hospitalizations

DHEC requires weekly submission of laboratory confirmed influenza hospitalizations. Hospitals must report these to their <u>regional</u> health department by noon on Monday for the preceding week.

Voluntary Networks

Influenza-Like Illness (ILINet) Sentinel Providers Network ILINet focuses on the number of patients presenting with influenza-like symptoms in the absence of another known cause. ILI is defined as fever (temperature $\geq 100^{\circ}$ F) plus a cough and/or a sore throat in the absence of another known cause. Providers submit weekly reports to the CDC of the total number of patients seen in a week and the subset number of those patients with ILI symptoms by age group.

South Carolina Disease Alerting, Reporting & Tracking System (SC-DARTS) (Syndromic surveillance)

SC-DARTS is a collaborative network of syndromic surveillance systems within South Carolina. Currently our network contains the following data sources: SC Hospital Emergency Department (ED) chief-complaint data, Poison Control Center call data, Over-the-Counter (OTC) pharmaceutical sales surveillance, and CDC's BioSense Biosurveillance system. The hospital ED syndromic surveillance system classifies ED chief complaint data into appropriate syndrome categories (ex: Respiratory, GI, Fever, etc.). These syndrome categories are then analyzed using the cumulative sum (CUSUM) methodology to detect any significant increases. Syndromic reports are distributed back to the hospital on a daily basis.

For additional information about SC-DARTS, contact the Syndromic Surveillance epidemiologist at cartere@dhec.sc.gov.

VII. Definitions for Influenza Surveillance

Activity level: Indicator of the geographic spread of influenza activity which is reported to CDC each week.

- *No activity:* No increase in ILI activity and no laboratory-confirmed influenza cases.
- Sporadic: No increase in ILI activity and isolated laboratory-confirmed influenza cases
- *Local:* Increased ILI or 2 or more institutional outbreaks in one region and laboratory-confirmed influenza cases within the past 3 weeks in the region with increased ILI or outbreaks
- *Regional:* Increased ILI or institutional outbreaks in 2-3 regions and laboratory-confirmed influenza cases within the past 3 weeks in the regions with increased ILI or institutional outbreaks
- *Widespread:* Increased ILI and/or institutional outbreaks in at least 4 regions and laboratory confirmed influenza in the state within the past 3 weeks

Confirmatory testing: Influenza testing which is considered to be confirmatory, such as a viral culture or RT-PCR

Fever-flu syndrome: Includes chief complaints with any of the following ICD codes or terms: flu, fev, high temp, temp10, feel hot, night sweat, FEB, shiver, FUO, chill, 780.6, 487, viral INF, pain all over, ILI, and body ache. Weekly fever flu count is the sum of all records, statewide, that were categorized into the fever flu syndrome. The state denominator is a broader modification of the respiratory syndrome that includes records that have fever flu chief complains and general respiratory illness complaints, which include: cough, coughing, URI, pneumonia, croup, bronchitis, and cold. The fever flu percentage equals (weekly fever flu count/weekly state denominator)*100.

Influenza-associated death: A death in which laboratory confirmation (see definition below) for influenza was reported, or for which an autopsy report consistent with influenza was provided, regardless of primary cause of death.

Influenza-like illness (ILI): Fever (temperature of 100°F [37.8°C] or greater) and cough and/or sore throat. The SC baseline is the mean percentage of patient visits for ILI during non-influenza weeks (weeks when percent of positive lab tests was below 20%) for the previous three seasons plus two standard deviations.

MMWR week: Term for influenza surveillance week. Each week begins on Sunday and ends on Monday. The influenza season begins with MMWR week 40 and ends with MMWR week 39. The 2015-16 influenza season began on October 4, 2015 and will end on October 1, 2016.

Laboratory-confirmation: Influenza positive resulting from one of the following methods:

- DFA
- IFA
- Rapid influenza antigen test
- RT-PCR
- Viral culture