



## This is an official CDC Health Advisory

Distributed via Health Alert Network  
December 31, 2021: 8:30 PM  
10509-CHA-12-31-2021-COVID-19-Therapeutics

### Using Therapeutics to Prevent and Treat COVID-19

#### Summary

The SARS-CoV-2 [Omicron](#) variant has quickly become the [dominant variant of concern](#) in the United States and is present in all 50 states. The Centers for Disease Control and Prevention (CDC) recommends that eligible individuals should get all [vaccines and booster shots](#) as the best preventive measure available against severe disease, hospitalizations, and death due to COVID-19.

Therapeutics are also available for preventing and treating COVID-19 in specific [at-risk populations](#). These therapeutics differ in efficacy, route of administration, risk profile, [and whether they are authorized by the U.S Food and Drug Administration \(FDA\) for adults only or adults and certain pediatric populations](#). Some therapeutics are in short supply, but availability is expected to increase in the coming months. This Health Alert Network (HAN) Health Advisory serves to familiarize healthcare providers with available therapeutics, understand how and when to prescribe [and prioritize](#) them, and recognize contraindications.

#### Background

On November 24, 2021, a new variant of SARS-CoV-2, B.1.1.529 (Omicron), was reported to the [World Health Organization](#) (WHO). On December 1, 2021, the first case of COVID-19 attributed to Omicron was reported in the United States. CDC has been working with state, tribal, local, and territorial public health officials to monitor the spread of the Omicron variant in the United States and has identified a [rapid increase in infections](#) consistent with what has been observed in other countries. Current [CDC recommendations for vaccines and booster shots](#) are expected to protect against severe illness, hospitalizations, and deaths from infection with the Omicron variant. Some studies have found lower effectiveness of the primary series of vaccines against infection and demonstrated the importance of booster doses (1-3). The United States Government is continuously working with private and public partners to bring new therapeutic options for use against SARS-CoV-2 variants of concern, including the Omicron variant.

### Monoclonal Antibodies

The Omicron variant, with its numerous mutations in the spike protein, is not neutralized by [bamlanivimab and etesevimab](#) or [casirivimab and imdevimab](#), the most frequently prescribed monoclonal antibody-based COVID-19 treatments (4-5). Despite some reduction in neutralization concentrations, [sotrovimab](#) remains effective against all variants of concern, including Omicron (6). However, sotrovimab is currently in limited supply, and [its use should be prioritized](#) for non-hospitalized patients with risk factors for progression to severe COVID-19, including individuals who are unvaccinated or have not received all [vaccines and booster shots as recommended by CDC](#), individuals with clinical risk factors, older age (for example  $\geq 65$  years of age), and [individuals not expected to mount an adequate immune response](#). Sotrovimab can be used in these [high-risk individuals](#) when Paxlovid (described below) is not indicated due to potential severe drug-drug interactions or if Paxlovid is not available.

### Antivirals

- [Remdesivir](#) is a nucleoside analog approved by FDA for the treatment of hospitalized patients with COVID-19. A recent randomized placebo-controlled outpatient study evaluated three daily intravenous (IV) infusion of remdesivir given within seven days of symptom onset. This study found that the reduction in hospitalization rates was similar to that achieved by using anti-SARS-CoV-2 monoclonal antibody-based therapy (7). Remdesivir is expected to be effective against the Omicron variant based on in vitro data; however, in vivo data are currently limited (8). Outpatient use of remdesivir requires support of IV infusion centers with appropriate skilled staffing.
- Two oral antivirals, [Paxlovid](#) (ritonavir-boosted nirmatrelvir) and [molnupiravir](#), are now available under Emergency Use Authorization by FDA for treating COVID-19 in outpatients with mild to moderate disease. Each drug is administered twice daily for five days. There are considerable differences in efficacy, risk profiles, and use restrictions between the two oral antivirals. From their individual clinical trials, compared to placebo, severe outcomes (hospitalization or death) were reduced by 88% for [Paxlovid](#) compared to 30% for molnupiravir (9). Healthcare providers need to be familiar with these distinctions to make clinical decisions and inform patients. In addition, initiating treatment with these oral antivirals must begin within five days of symptom onset to maintain product efficacy.
  - [Paxlovid](#) is currently in very limited supply and use should be prioritized for [higher risk populations](#). Due to the potential for severe drug-drug interactions with ritonavir, a medication used for HIV treatment, CDC strongly suggests that healthcare providers not experienced in prescribing [Paxlovid](#) refer to the [NIH Statement on Paxlovid Drug-Drug Interactions | COVID-19 Treatment Guidelines](#). Healthcare providers could also contact a local clinical pharmacist or an infectious disease specialist for advice.
  - Like Paxlovid, molnupiravir is expected to be active against all circulating variants of concern, including Omicron (8). Molnupiravir should only be used when other options are not available, due to its lower efficacy. [Molnupiravir use is not recommended](#) in pregnancy because of potential mutagenicity.

[Molnupiravir is also not recommended](#) in patients who are breastfeeding or pediatric patients due to limited data within these populations and concerns for potential bone growth toxicity in young individuals.

#### *Pre-exposure therapeutics for high-risk groups*

AstraZeneca's [EVUSHELD](#), which includes two long-acting anti-SARS-CoV-2 monoclonal antibodies, is the only Emergency Use Authorization pre-exposure prophylaxis product available. EVUSHELD is expected to be effective against the Omicron variant; however, treatment effectiveness should be monitored. EVUSHELD is intended for the highest risk immunocompromised patients who are not expected to have an effective response to vaccination. EVUSHELD is indicated for pre-exposure prophylaxis only and not for treatment of patients with COVID-19.

### **Recommendations for Healthcare Providers**

- As with all therapeutics, the best use of therapeutics includes an appropriate clinical assessment and an up-to-date and informed risk-benefit discussion to address any questions or concerns from patients.
- Obtain further information on clinical use of products through [NIH's COVID-19 Treatment Guidelines](#), the [Assistant Secretary for Preparedness and Response Public Health Emergency COVID-19 Therapeutics site](#), and through professional societies such as [IDSA's Guidelines on the Management of Patients with COVID-19](#).
- Check with state and local health departments on key sites that have been identified for distribution of therapeutics, including cancer treatment centers and oncology providers.
  - Monoclonal Antibody Treatment Sites in South Carolina can be found on the [DHEC website](#).
- If the Delta variant still represents a significant proportion of infections in a region and other options are not available or are contraindicated, eligible patients can be offered [bamlanivimab and etesevimab](#) or [casirivimab and imdevimab](#), with the understanding that these treatments would be ineffective against the Omicron variant. This concern can be mitigated if [virus-specific diagnostic testing](#) in a given patient indicates infection with the Omicron variant is unlikely.
- Prioritize high risk patients, particularly if therapeutics are in short supply, using [NIH COVID-19 Treatment Guidelines when supply constraints exist](#). This document presents a tiered approach to prioritization.
- Continue to encourage COVID-19 vaccination, including booster vaccination.

### **For More Information**

- [Omicron Variant: What You Need to Know | CDC](#)
- [Interim Clinical Considerations for Use of COVID-19 Vaccines | CDC](#)
- [CDC COVID Data Tracker](#)
- [COVID-19 Treatment Guidelines: What's New](#)

- [COVID-19 Treatment Guidelines: Antiviral Therapy](#)
- [NIH Statement on Therapies for High-Risk, Nonhospitalized Patients | COVID-19 Treatment Guidelines](#)
- [NIH Statement on Paxlovid Drug-Drug Interactions | COVID-19 Treatment Guidelines](#)
- [The COVID-19 Treatment Guidelines Panel's Interim Statement on Patient Prioritization for Outpatient Anti-SARS-CoV-2 Therapies or Preventive Strategies When There Are Logistical or Supply Constraints](#)
- [Side by Side Overview of Outpatient Therapies Authorized for Treatment of Mild-Moderate COVID-19](#)

## References

1. Andrews N, Stowe J, Kirseborm F, et al. Effectiveness of COVID-19 vaccines against the Omicron (B.1.1529) variant of concern. medRxiv 2021. doi: <https://doi.org/10.1101/2021.12.14.21267615>
2. Ferguson N, Ghani A, Cori A, et al. Report 49: Growth, population distribution and immune escape of Omicron in England. Imperial College London (2021-12-20). doi: <https://doi.org/10.25561/93038> (direct link: <https://spiral.imperial.ac.uk/handle/10044/1/93038>)
3. Discovery Health. Real world analysis of Omicron outbreak in South Africa including vaccine effectiveness. Accessed 2021-12-23 at <https://www.discovery.co.za/corporate/news-room> (direct link: [https://resources.mynewsdesk.com/image/upload/fl\\_attachment/lw9szzdtqfwwitkfbcoq](https://resources.mynewsdesk.com/image/upload/fl_attachment/lw9szzdtqfwwitkfbcoq))
4. Aggarwal A, Ospina Stella A, Walker G, et al. SARS-CoV-2 Omicron: evasion of potent humoral responses and resistance to clinical immunotherapeutics relative to viral variants of concern. medRxiv 2021. doi: <https://doi.org/10.1101/2021.12.14.21267772>
5. Planas D, Saunders N, Maes P, et al. Considerable escape of SARS-CoV-2 variant Omicron to antibody neutralization. bioRxiv 2021. doi: <https://doi.org/10.1101/2021.12.14.472630>
6. Cathcart A, Havenar-Daughton C, Lempp F, et al. The dual function monoclonal antibodies VIR7831 and VIR-7832 demonstrate potent in vitro and in vivo activity against SARS-CoV-2. bioRxiv 2021. doi: <https://doi.org/10.1101/2021.03.09.434607>
7. Gottlieb, Robert L et al. "Early Remdesivir to Prevent Progression to Severe Covid-19 in Outpatients." The New England Journal of Medicine (2021): <https://www.nejm.org/doi/full/10.1056/NEJMoa2116846>
8. Vangeel L, De Jonghe S, Maes P, et al. Remdesivir, Molnupiravir and Nirmatrelvir remain active against SARS-CoV-2 Omicron and other variants of concern. bioRxiv 2021. doi: <https://doi.org/10.1101/2021.12.27.474275>
9. Bernal, A, Gomes da Silva, M, Musungaie, D, et al. Molnupiravir for Oral Treatment of Covid-19 in Nonhospitalized Patients. The New England Journal of Medicine (2021). doi: [10.1056/NEJMoa2116044](https://doi.org/10.1056/NEJMoa2116044)

## DHEC contact information for reportable diseases and reporting requirements

Reporting of **SARS-CoV-2 laboratory results (positive, negative, and indeterminate), whole genome sequencing results, and COVID-19 associated deaths** is consistent with South Carolina

Law requiring the reporting of diseases and conditions to your state or local public health department. (State Law # 44-29-10 and Regulation # 61-20) as per the DHEC 2021 List of Reportable Conditions available at:

<https://www.scdhec.gov/sites/default/files/Library/CR-009025.pdf>

Federal HIPAA legislation allows disclosure of protected health information, without consent of the individual, to public health authorities to collect and receive such information for the purpose of preventing or controlling disease. (HIPAA 45 CFR §164.512).

<b>Regional Public Health Offices – 2021</b>			
Mail or call reports to the Epidemiology Office in each Public Health Region			
<b>MAIL TO:</b>			
<b><u>Lowcountry</u></b> 4050 Bridge View Drive, Suite 600 N. Charleston, SC 29405 Fax: (843) 953-0051	<b><u>Midlands</u></b> 2000 Hampton Street Columbia, SC 29204 Fax: (803) 576-2993	<b><u>Pee Dee</u></b> 1931 Industrial Park Road Conway, SC 29526 Fax: (843) 915-6506	<b><u>Upstate</u></b> 200 University Ridge Greenville, SC 29602 Fax: (864) 282-4373
<b>CALL TO:</b>			
<b><u>Lowcountry</u></b> Allendale, Bamberg, Beaufort, Berkeley, Calhoun, Charleston, Colleton, Dorchester, Hampton, Jasper, Orangeburg  Office: (843) 441-1091 Nights/Weekends: (843) 441-1091	<b><u>Midlands</u></b> Aiken, Barnwell, Chester, Edgefield, Fairfield, Kershaw, Lancaster, Lexington, Newberry, Richland, Saluda, York  Office: (888) 801-1046 Nights/Weekends: (888) 801-1046	<b><u>Pee Dee</u></b> Clarendon, Chesterfield, Darlington, Dillon, Florence, Georgetown, Horry, Lee, Marion, Marlboro, Sumter, Williamsburg  Office: (843) 915-8886 Nights/Weekends: (843) 915-8845	<b><u>Upstate</u></b> Abbeville, Anderson, Cherokee, Greenville, Greenwood, Laurens, McCormick, Oconee, Pickens, Spartanburg, Union  Office: (864) 372-3133 Nights/Weekends: (864) 423-6648
<b>For information on reportable conditions, see</b> <a href="https://www.scdhec.gov/ReportableConditions">https://www.scdhec.gov/ReportableConditions</a>		<b><u>DHEC Bureau of Communicable Disease Prevention &amp; Control</u></b> <b>Division of Acute Disease Epidemiology</b> 2100 Bull St • Columbia, SC 29201 Phone: (803) 898-0861 • Fax: (803) 898-0897 Nights / Weekends: 1-888-847-0902	

Categories of Health Alert messages:

- Health Alert** Conveys the highest level of importance; warrants immediate action or attention.
- Health Advisory** Provides important information for a specific incident or situation; may not require immediate action.
- Health Update** Provides updated information regarding an incident or situation; unlikely to require immediate action.
- Info Service** Provides general information that is not necessarily considered to be of an emergent nature.