



## This is an official **CDC Health Update**

Distributed via Health Alert Network  
May 25, 2016; 5:00 PM  
10379-CHU-05-25-2016-ZIKA

### **Diagnostic Testing of Urine Specimens for Suspected Zika Virus Infection**

#### **Summary**

On May 13, 2016, the Centers for Disease Control and Prevention (CDC) issued [interim guidance](http://www.cdc.gov/mmwr/volumes/65/wr/mm6518e1.htm) (<http://www.cdc.gov/mmwr/volumes/65/wr/mm6518e1.htm>) that recommends Zika virus rRT-PCR testing of urine collected less than 14 days after symptom onset, along with testing of patient-matched serum samples, for the diagnosis of suspected Zika virus infection (1). The purpose of this Health Alert Network (HAN) health update is to further disseminate information about the interim guidance to clinical and public health professionals.

#### **Background**

Zika virus is a mosquito-borne flavivirus. Zika virus infection during pregnancy can cause microcephaly and other severe fetal brain defects. Zika virus infection is also associated with Guillain-Barré syndrome. Transmission of Zika can occur through mosquito bite, from a pregnant woman to her fetus, through sexual contact with an infected male, and possibly through blood transfusion. The most common symptoms of Zika virus disease are fever, rash, joint pain, or conjunctivitis. Other common symptoms include muscle pain and headache. Evidence from case reports and experience from related flavivirus infections indicate that the incubation period for Zika is likely a few to 14 days.

Diagnostic testing for Zika virus infection can be accomplished using molecular and serologic methods. The U.S. Food and Drug Administration (FDA) has issued [Emergency Use Authorizations](http://www.fda.gov/MedicalDevices/Safety/EmergencySituations/ucm161496.htm) (EUA) (<http://www.fda.gov/MedicalDevices/Safety/EmergencySituations/ucm161496.htm>) for several diagnostic assays to detect Zika virus infection (2). The EUAs authorize real-time reverse transcription-polymerase chain reaction (rRT-PCR) assays to detect Zika virus RNA in specified clinical sample types, and an immunoglobulin M (IgM) antibody capture enzyme-linked immunosorbent assay (ELISA) to detect anti-Zika virus IgM antibodies in serum and cerebrospinal fluid. The CDC Trioplex rRT-PCR assay is authorized by FDA for Zika virus testing of urine and serum. Anti-Zika IgM antibodies develop during the first week of illness and persist for approximately 12 weeks following infection. However, extensive cross-reactivity can occur in flavivirus serological assays, and therefore additional tests, such as the plaque reduction neutralization test (PRNT), are necessary to distinguish Zika virus infection from other flavivirus infections.

Although Zika virus RNA is unlikely to be detected in serum after the first week of illness, recent data suggest that Zika virus RNA can persist in urine for at least two weeks post symptom onset (3). Given this information, on May 13, 2016, CDC issued interim guidance on rRT-PCR testing for Zika virus RNA in urine (1). CDC now recommends that, for persons with suspected Zika virus disease, Zika virus rRT-PCR should be performed on both urine and serum specimens collected within 7 days after onset of symptoms. Zika virus rRT-PCR also should be performed on urine specimens collected within 14 days after onset of symptoms. A positive rRT-PCR result in either specimen confirms Zika virus infection. However, a negative rRT-PCR in a serum or urine sample collected at any time point after illness onset

does not exclude Zika virus infection, and in these cases IgM antibody testing should be performed on serum.

CDC recommendations for Zika virus testing of serum and other clinical specimens remain unchanged at this time. Please contact your state or local health department to facilitate testing.

### **Recommendations for Health Care Providers and Public Health Practitioners**

- Collect urine samples within 14 days post symptom onset along with patient-matched serum samples for those who match CDC Zika virus clinical and/or epidemiological testing criteria for Zika virus infection.
- Perform Zika virus rRT-PCR testing on urine, in conjunction with testing of serum using the appropriate molecular or serologic assay, based on days post symptom onset.

### **Additional Considerations**

- Further investigation is needed to determine the sensitivity and utility of Zika virus rRT-PCR on urine specimens collected  $\geq 14$  days after onset of symptoms: limited data in pregnant women suggest that viremia in serum might be prolonged in pregnancy (4, 5).

### **References**

1. CDC. Interim guidance for Zika virus testing of Urine – United States, 2016. MMWR Morb Mortal Wkly Rep 2016; 65. DOI: <http://dx.doi.org/10.15585/mmwr.mm6518e1>.
2. Food and Drug Administration. Emergency Use Authorizations. <http://www.fda.gov/MedicalDevices/Safety/EmergencySituations/ucm161496.htm>
3. Comparison of Test Results for Zika Virus RNA in Urine, Serum, and Saliva Specimens from Persons with Travel-Associated Zika Virus Disease — Florida, 2016 <http://www.cdc.gov/mmwr/volumes/65/wr/mm6518e2.htm>
4. Driggers RW, Ho CY, Korhonen EM, et al. Zika virus infection with prolonged maternal viremia and fetal brain abnormalities. N Engl J Med. March 30, 2016. DOI: 10.1056/NEJMoa1601824
5. Bocanegra C. Zika virus infection in pregnant women in Barcelona, Spain. Clin Microbiol Infect. April 3, 2016. DOI: 10.1016/j.cmi.2016.03.025.

### **For More Information**

- General information about Zika virus and disease: <http://www.cdc.gov/zika/>
- Zika virus information for clinicians: <http://www.cdc.gov/zika/hc-providers/index.html>
- Memorandum – Revised diagnostic testing for Zika, chikungunya, and dengue viruses in US Public Health Laboratories (not updated with urine guidance): <http://www.cdc.gov/zika/state-labs/index.html>
- Diagnostic testing: <http://www.cdc.gov/zika/hc-providers/diagnostic.html>
- Collection and submission of body fluids for Zika virus testing: <http://www.cdc.gov/zika/hc-providers/body-fluids-collection-submission.html>

*The Centers for Disease Control and Prevention (CDC) protects people's health and safety by preventing and controlling diseases and injuries; enhances health decisions by providing credible information on critical health issues; and promotes healthy living through strong partnerships with local, national, and international organizations.*

---

### **Categories of Health Alert Network messages:**

**Health Alert** Requires immediate action or attention; highest level of importance

**Health Advisory** May not require immediate action; provides important information for a specific incident or situation

**Health Update** Unlikely to require immediate action; provides updated information regarding an incident or situation

**HAN Info Service** Does not require immediate action; provides general public health information

## DHEC contact information for reportable diseases and reporting requirements

Reporting of Zika Virus is consistent with South Carolina Law requiring the reporting of diseases and conditions to your state or local public health department. (State Law # 44-29-10 and Regulation # 61-20) as per the DHEC 2016 List of Reportable Conditions available at:

<http://www.scdhec.gov/Library/CR-009025.pdf>

Federal HIPAA legislation allows disclosure of protected health information, without consent of the individual, to public health authorities to collect and receive such information for the purpose of preventing or controlling disease. (HIPAA 45 CFR §164.512).

<b>Regional Public Health Offices – 2016</b>			
Mail or call reports to the Epidemiology Office in each Public Health Region			
<b>MAIL TO:</b>			
<b>Lowcountry</b> 4050 Bridge View Drive, Suite 600 N. Charleston, SC 29405 Fax: (843) 953-0051	<b>Midlands</b> 2000 Hampton Street Columbia, SC 29204 Fax: (803) 576-2993	<b>Pee Dee</b> 145 E. Cheves Street Florence, SC 29506 Fax: (843) 661-4859	<b>Upstate</b> 200 University Ridge Greenville, SC 29602 Fax: (864) 282-4373
<b>CALL TO:</b>			
<b>Lowcountry</b> <b>Berkeley, Charleston, Dorchester</b> Phone: (843) 953-0043 Nights/Weekends: (843) 441-1091  <b>Beaufort, Colleton, Hampton, Jasper</b> Phone: (843) 322-2453 Nights/Weekends: (843) 441-1091  <b>Allendale, Bamberg, Calhoun, Orangeburg</b> Phone: (803) 268-5833 Nights/Weekends: (843) 441-1091	<b>Midlands</b> <b>Kershaw, Lexington, Newberry, Richland</b> Phone: (803) 576-2749 Nights/Weekends: (888) 801-1046  <b>Chester, Fairfield, Lancaster, York</b> Phone: (803) 286-9948 Nights/Weekends: (888) 801-1046  <b>Aiken, Barnwell, Edgefield, Saluda</b> Phone: (803) 642-1618 Nights/Weekends: (888) 801-1046	<b>Pee Dee</b> <b>Chesterfield, Darlington, Dillon, Florence, Marlboro, Marion</b> Phone: (843) 661-4830 Nights/Weekends: (843) 915-8845  <b>Clarendon, Lee, Sumter</b> Phone: (803) 773-5511 Nights/Weekends: (843) 915-8845  <b>Georgetown, Horry, Williamsburg</b> Phone: (843) 915-8804 Nights/Weekends: (843) 915-8845	<b>Upstate</b> <b>Anderson, Oconee</b> Phone: (864) 260-5801 Nights/Weekends: (866) 298-4442  <b>Abbeville, Greenwood, Laurens, McCormick</b> Phone: (864) 227-5947 Nights/Weekends: (866) 298-4442  <b>Cherokee, Greenville, Pickens, Spartanburg, Union</b> Phone: (864) 372-3133 Nights/Weekends: (866) 298-4442
<b>For information on reportable conditions, see</b> <b><a href="http://www.scdhec.gov/Health/FHPF/ReportDiseaseAdverseEvents/ReportableConditionsInSC/">http://www.scdhec.gov/Health/FHPF/ReportDiseaseAdverseEvents/ReportableConditionsInSC/</a></b>		<b>DHEC Bureau of Disease Control</b> <b>Division of Acute Disease Epidemiology</b> 2100 Bull St • Columbia, SC 29201 Phone: (803) 898-0861 • Fax: (803) 898-0897 Nights / Weekends: 1-888-847-0902	

Categories of Health Alert messages:

<b>Health Alert</b>	Conveys the highest level of importance; warrants immediate action or attention.
<b>Health Advisory</b>	Provides important information for a specific incident or situation; may not require immediate action.
<b>Health Update</b>	Provides updated information regarding an incident or situation; unlikely to require immediate action.
<b>Info Service</b>	Provides general information that is not necessarily considered to be of an emergent nature.